

brought to you by [www.thebacteriophages.org](#) and [www.phage.org](#)

REFERENCES

1. **Abril, A.M., M. Salas, and J.M. Hermoso.** 2000. Identification of residues within two regions involved in self-association of viral histone-like protein p6 from phage ϕ 29. *J. Biol. Chem.* **275**:26404-26410.
2. **Anderson, D.L., and B.E. Reilly.** 1993. Morphogenesis of bacteriophage ϕ 29, in: *Bacillus subtilis and other gram-positive bacteria: biochemistry, physiology, and molecular genetics*. L. Sonenshein, J.A. Hoich, and R. Losick, (eds.), pp. 859-867, American Society for Microbiology, Washington, D.C.
3. **Barthelemy, I., R.P. Mellado, and M. Salas.** 1989. *In vitro* transcription of bacteriophage ϕ 29 DNA: Inhibition of early promoters by the viral replication protein p6. *J. Virol.* **63**:460-462.
4. **Barthelemy, I., M. Salas, and R.P. Mellado.** 1987. *In vivo* transcription of bacteriophage ϕ 29 DNA. Transcription termination. *J. Virol.* **61**:1751-1755.
5. **Benes, V., L. Arnold, J. Smrt, and V. Paces.** 1989. Nucleotide sequence of the right early region of Bacillus phage ϕ 15 and comparison with related phages: reorganization of gene 17 during evolution. *Gene* **75**:341-347.
6. **Bernad, A., L. Blanco, J.M. Lázaro, G. Martín, and M. Salas.** 1989. A conserved 3'→ 5' exonuclease active site in prokaryotic and eukaryotic DNA polymerases. *Cell* **59**:219-228.
7. **Bernad, A., J.M. Lázaro, M. Salas, and L. Blanco.** 1990. The highly conserved amino acid sequence motif Tyr-Gly-Asp-Thr-Asp-Ser in α -like DNA polymerases is required by phage ϕ 29 DNA polymerase for protein-primed initiation and polymerization. *Proc. Natl. Acad. Sci. USA* **87**:4610-4614.
8. **Bjornsti, M.A., B.E. Reilly, and D.L. Anderson.** 1981. *In vitro* assembly of the *Bacillus subtilis* bacteriophage ϕ 29. *Proc. Natl. Acad. Sci. USA* **78**:5861-5865.
9. **Bjornsti, M.A., B.E. Reilly, and D.L. Anderson.** 1983. Morphogenesis of bacteriophage ϕ 29 of *Bacillus subtilis*: oriented and quantized *in vitro* packaging of DNA protein gp3. *J. Virol.* **45**:383-396.

10. **Blanco, L., A. Bernad, J.M. Lázaro, G. Martín, C. Garmendia, and M. Salas.** 1989. Highly efficient DNA synthesis by the phage ϕ 29 DNA polymerase. Symmetrical mode of DNA replication. *J. Biol. Chem.* **264**:8935-8940.
11. **Blanco, L. J. Gutiérrez, J.M. Lázaro, A. Bernad, and M. Salas.** 1986. Replication of phage ϕ 29 DNA *in vitro*: role of the viral protein p6 in initiation and elongation. *Nucleic Acids Res.* **14**:4923-4937.
12. **Blanco, L., J.M. Lázaro, M. de Vega, A. Bonnin, and M. Salas.** 1994. Terminal protein-primed DNA amplification. *Proc. Natl. Acad. Sci. USA* **91**:12198-12202.
13. **Blanco, L., I. Prieto, J. Gutiérrez, A. Bernad, J.M. Lázaro, J.M. Hermoso, and M. Salas.** 1987. Effect of NH_4^+ ions on ϕ 29 DNA-protein p3 replication: Formation of a complex between the terminal protein and the DNA polymerase. *J. Virol.* **61**:3983-3991.
14. **Blanco, L., and M. Salas.** 1984. Characterization and purification of a phage ϕ 29 coded DNA polymerase required for the initiation of replication. *Proc. Natl. Acad. Sci. USA* **81**:5325-5329.
15. **Blanco, L., and M. Salas.** 1985a. Replication of ϕ 29 DNA with purified terminal protein and DNA polymerase: synthesis of full-length ϕ 29 DNA. *Proc. Natl. Acad. Sci. USA* **82**:6404-6408.
16. **Blanco, L., and M. Salas.** 1985b. Characterization of a 3'→ 5' exonuclease activity in the phage ϕ 29-encoded DNA polymerase. *Nucl. Acids Res.* **13**:1239-1249.
17. **Blanco, L., and M. Salas.** 1995. Mutational analysis of bacteriophage ϕ 29 DNA polymerase. *Methods Enzymol.* **262**:283-294.
18. **Blanco, L., and M. Salas.** 1996. Relating structure to function in ϕ 29 DNA polymerase. *J. Biol. Chem.* **271**:8509-8512.
19. **Blasco, M.A., A. Bernad, L. Blanco, and M. Salas.** 1991. Characterization and mapping of the pyrophosphorolytic activity of the phage ϕ 29 DNA polymerase. *J. Biol. Chem.* **266**:7904-7909.

20. **Blasco, M.A., J.M. Lázaro, A. Bernad, L. Blanco, and M. Salas.** 1992. ϕ 29 DNA polymerase active site: mutants in conserved residues Tyr254 and Tyr390 are affected in dNTP binding. *J. Biol. Chem.* **267**:19427-19434.
21. **Blasco, M.A., J.M. Lázaro, L. Blanco and M. Salas.** 1993a. ϕ 29 DNA polymerase active site. Residue Asp²⁴⁹ of conserved amino acid motif Dx₂SLYP is critical for synthetic activities. *J. Biol. Chem.* **268**:24106-24113.
22. **Blasco, M.A., J.M. Lázaro, L. Blanco, and M. Salas.** 1993b. ϕ 29 DNA polymerase active site. The conserved amino acid motif "Kx₃NSxYG" is involved in template-primer binding and dNTP selection. *J. Biol. Chem.* **268**:16763-16770.
23. **Blasco, M.A., J. Méndez, J.M. Lázaro, L. Blanco, and M. Salas.** 1995. Primer-terminus stabilization at the ϕ 29 DNA polymerase active site: mutational analysis of conserved motif KxY. *J. Biol. Chem.* **270**:2735-2740.
24. **Bonnin, A., J.M. Lázaro, L. Blanco, and M., Salas.** 1999. A single tyrosine prevents insertion of ribonucleotides in the eukaryotic-type ϕ 29 DNA polymerase. *J.Mol.Biol.* **290**:241-251.
25. **Bravo, A., and M. Salas.** 1997. Initiation of bacteriophage ϕ 29 DNA replication *in vivo*: Assembly of a membrane-associated multiprotein complex. *J. Mol. Biol.* **269**:102-112.
26. **Bravo, A., and M. Salas.** 1998. Polymerization of bacteriophage ϕ 29 replication protein p1 into protofilament sheets. *EMBO J.* **17**:6096-6105.
27. **Bravo, A., B. Illana, and M. Salas.** 2000. Compartmentalization of phage ϕ 29 DNA replication: Interaction between the primer terminal protein and the membrane-associated protein p1. *EMBO J.* **19**:5575-5584.
28. **Caldentey, J., L. Blanco, D.H. Bamford, and M. Salas.** 1993. *In vitro* replication of bacteriophage PRD1 DNA. Characterization of the protein-primed initiation site. *Nucl. Acids Res.* **21**:3725-3730.
29. **Calles, B., M. Salas, and F. Rojo.** 2002. The ϕ 29 transcriptional regulator contacts the nucleoid protein p6 to organize a repression complex. *EMBO J.* **21**:6185-6194.

30. **Camacho, A., and M. Salas.** 1999. Effect of mutations in the "Extended -10" motif of three *Bacillus subtilis* σ^A -RNA polymerase-dependent promoters. *J.Mol.Biol.* **286**:683-693.
31. **Camacho, A., and M. Salas.** 2001a. Repression of bacteriophage ϕ 29 early promoter C2 by the viral protein p6 is due to the impairment of the closed complex. *J.Biol.Chem.* **276**: 28927-28932.
32. **Camacho, A., and M. Salas.** 2001b. Mechanism for the switch of ϕ 29 DNA early to late transcription by regulatory protein p4 and histone-like protein p6. *EMBO J.* **20**:6060-6070.
33. **Camacho, A., F. Jiménez, J. de la Torre, J.L. Carrascosa, R.P. Mellado, C. Vásquez, E. Viñuela, and M. Salas.** 1977. Assembly of *B. subtilis* phage ϕ 29. I. Mutants in the cistrons coding for the structural protein. *Eur. J. Biochem.* **73**:39-55.
34. **Camacho, A., F., Jiménez, E. Viñuela, and M. Salas.** 1979. Order of assembly of the lower collar and the tail proteins of *Bacillus subtilis* phage ϕ 29. *J. Virol.* **29**:540-545.
35. **Carazo, J.M., L.E. Donate, L. Herranz, J.P. Secilla, and J.L. Carrascosa.** 1986. Three-dimensional reconstruction of the connector of bacteriophage ϕ 29 at 1.8 nm resolution. *J.Mol.Biol.* **192**:853-867.
36. **Carrascosa, J.L., A. Camacho, F. Moreno, F. Jiménez, R.P. Mellado, E., Viñuela, and M. Salas.** 1976. *Bacillus subtilis* phage ϕ 29: Characterization of gene products and functions. *Eur. J. Biochem.* **66**:229-241.
37. **Carrascosa, J.L., J.M. Carazo, and N. García.** 1983. Structural localization of the proteins of the head to tail connecting region of bacteriophage ϕ 29. *Virology* **124**:133-144.
38. **Crucitti, P., A.M. Abril, and M. Salas.** 2003. Bacteriophage ϕ 29 early protein p17. Self-association and hetero-association with the viral histone-like protein p6. *J.Biol.Chem.* **278**:4906-4911.

39. **Crucitti, P., J.M. Lázaro, V. Benes, and M. Salas.** 1998. Bacteriophage ø29 early protein p17 is conditionally required for the first rounds of viral DNA replication. *Gene* **223**:135-142.
40. **de Vega, M., L. Blanco, and M. Salas.** 1998a. ø29 DNA polymerase residue Ser¹²², a ssDNA ligand for 3'-5' exonucleolysis, is required to interact with the terminal protein. *J. Biol. Chem.* **273**:28966-28977.
41. **de Vega, M., T. Ilyina, J.M. Lázaro, M. Salas, and L. Blanco.** 1997. An invariant lysine residue is involved in catalysis at the 3'-5' exonuclease active site of eukaryotic-type DNA polymerases. *J. Mol. Biol.* **270**:65-78.
42. **de Vega, M., J.M. Lázaro, M. Salas, and L. Blanco.** 1996. Primer-terminus stabilization at the 3'-5' exonuclease active site of ø29 DNA polymerase. Involvement of two amino acid residues highly conserved in proofreading DNA polymerases. *EMBO J.* **15**:1182-1192.
43. **de Vega, M., J.M. Lázaro, M. Salas, and L. Blanco.** 1998b. Mutational analysis of ø29 DNA polymerase residues acting as ssDNA ligands for 3'-5' exonucleolysis. *J. Mol. Biol.* **279**:807-822.
44. **Dufour, E., J. Méndez, J.M. Lázaro, M. de Vega, L. Blanco, and M., Salas.** 2000. An aspartic acid residue in TPR-1, a specific region of protein-priming DNA polymerases, is required for the functional interaction with primer terminal protein. *J.Mol.Biol.* **304**:289-300.
45. **Elías-Arnanz, M., and M. Salas.** 1999. Functional interactions between a phage histone-like protein and a transcriptional factor in regulation of ø29 early-late transcriptional switch. *Genes Dev.* **13**:2502-2513.
46. **Escarmís, C., D. Guirao, and M. Salas.** 1989. Replication of recombinant ø29 DNA molecules in *Bacillus subtilis* protoplasts. *Virology* **169**:152-160.
47. **Esteban, J.A., L. Blanco, L. Villar, and M. Salas.** 1997. *In vitro* evolution of terminal protein-containing genomes. *Proc. Natl. Acad. Sci. USA* **94**:2921-2926.

48. **Esteban, J.A., M. Salas, and L. Blanco.** 1993. Fidelity of ϕ 29 DNA polymerase. Comparison between protein-primed initiation and DNA polymerization. *J. Biol. Chem.* **268**:2719-2726.
49. **Esteban, J.A., M.S. Soengas, M. Salas and L. Blanco.** 1994. 3'→5' exonuclease active site of ϕ 29 DNA polymerase. Evidence favoring a metal-ion-assisted reaction mechanism. *J. Biol. Chem.* **269**:31946-31954.
50. **Freire, R., M. Salas, and J.M. Hermoso.** 1994. A new protein domain for binding to DNA through the minor groove. *EMBO J.* **13**:4353-4360.
51. **Freire, R., M. Serrano, M. Salas, and J.M. Hermoso.** 1996. Activation of replication origins in ϕ 29-related phages requires the recognition of initiation proteins to specific nucleoprotein complexes. *J. Biol. Chem.* **271**:31000-31007.
52. **García, J.A., J.L. Carrascosa, and M. Salas.** 1983. Assembly of the tail protein of the *Bacillus subtilis* phage ϕ 29. *Virology* **12**:18-30.
53. **García, J.A., M.A. Peñalva, L. Blanco, and M. Salas.** 1984. Template requirements for initiation of phage ϕ 29 DNA replication *in vitro*. *Proc. Natl. Acad. Sci. USA* **81**:80-84.
54. **Garmendia, C., A. Bernad, J.A. Esteban, L. Blanco, and M. Salas.** 1992. The bacteriophage ϕ 29 DNA polymerase, a proofreading enzyme. *J. Biol. Chem.* **267**:2594-2599.
55. **Garmendia, C., M. Salas, and J.M. Hermoso.** 1988. Site-directed mutagenesis of bacteriophage ϕ 29 terminal protein: isolation and characterization of a Ser₂₃₂→thr mutant in the linking site. *Nucl. Acids Res.* **16**:5727-5740.
56. **Gascón, I., J.L. Carrascosa, L. Villar, J.M., Lázaro, and M. Salas.** 2002. Importance of the N-terminal region of the phage GA-1 SSB for its self-interaction ability and functionality. *J.Biol.Chem.* **277**:22534-22540.
57. **Gascón, I., C. Gutiérrez, and M. Salas.** 2000a. Structural and functional comparative study of the complexes formed by viral ϕ 29, Nf and GA-1 SSB proteins with DNA. *J.Mol.Biol.* **296**:989-999.

58. **Gascón, I., J.M. Lázaro, and M. Salas.** 2000b. Comparative functional characterization of viral ø29, Nf and GA-1 SSB proteins. *Nucl. Acids Res.* **28**:2034-2042.
59. **González-Huici, V., J.M. Lázaro, M. Salas, and J.M. Hermoso.** 2000a. Specific recognition of parental terminal protein by DNA polymerase for initiation of protein-primed DNA replication. *J. Biol.Chem.* **275**:14678-14683.
60. **González-Huici, V., M. Salas, and J.M. Hermoso.** 2000b. Sequence requirements for protein-primed initiation and elongation of phage ø29 DNA replication. *J.Biol.Chem.* **275**:14678-14683.
61. **Grimes, S., and D. Anderson.** 1990. RNA dependence of the bacteriophage ø29 DNA packaging ATPase. *J.Mol.Biol.* **215**:559-566.
62. **Guasch, A., J. Pous, A. Párraga, J.M. Valpuesta, J.M. Carrascosa, and M. Coll.** 1998. Crystallographic analysis reveals the 12-fold symmetry of the bacteriophage ø29 connector particles. *J.Mol.Biol.* **281**:219-225.
63. **Guo, P., S. Erickson, W. Xu, N. Olson, T.S. Baker, and D. Anderson.** 1991. Regulation of the phage ø29 prohead shape and size by the portal vertex. *Virology* **183**:366-373.
64. **Guo, P., S. Grimes, and D. Anderson.** 1986. A defined system for *in vitro* packaging of DNA-gp3 of the *Bacillus subtilis* bacteriophages ø29. *Proc.Natl.Acad.Sci.USA* **83**:3505-3509.
65. **Guo, P., C. Peterson, and D. Anderson.** 1987a. Initiation events in *in vitro* packaging of bacteriophage ø29 DNA-p3. *J.Mol.Biol.* **197**:219-228.
66. **Guo, P., C. Peterson, and D. Anderson.** 1987b. Prohead and DNA-gp3-dependent ATPase activity of the DNA packaging protein gp16 of bacteriophage ø29. *J.Mol.Biol.* **197**:229-236.
67. **Guo, P., C. Zhang, C. Chen, K. Garver, and M. Trottier.** 1998. Inter-RNA interaction of phage ø29 pRNA to form a hexameric complex for viral DNA transportation. *Mol.Cell.* **2**:149-155.

68. **Gutiérrez, C., R. Freire, M. Salas, and J.M. Hermoso.** 1994. Assembly of phage ø29 genome with viral protein p6 into a compact complex. *EMBO J.* **13**:269-276.
69. **Gutiérrez, C., G. Martín, J.M. Sogo, and M. Salas.** 1991. Mechanism of stimulation of DNA replication by bacteriophage ø29 SSB protein p5. *J. Biol. Chem.* **266**:2104-2111.
70. **Gutiérrez, J., J.A. García, L. Blanco, and M. Salas.** 1986a. Cloning and template activity of the origins of replication of phage ø29 DNA. *Gene* **43**:1-11.
71. **Gutiérrez, J., J. Vinós, I. Prieto, E. Méndez, J.M. Hermoso, and M. Salas.** 1986b. Signals in the ø29 DNA-terminal protein template for the initiation of phage ø29 DNA replication. *Virology* **155**:474-483.
72. **Hagen, E.W., B.E. Reilly, M.E. Tosi, and D.L. Anderson.** 1976. Analysis of gene function of bacteriophage ø29 of *Bacillus subtilis*: identification of cistrons essential for viral assembly. *J. Virol.* **19**:501-517.
73. **Harding, N.E., and J. Ito.** 1980. DNA replication of bacteriophage ø29: Characterization of the intermediates and location of the termini of replication. *Virology* **104**:323-338.
74. **Hendrix, R.W.** 1998. Bacteriophage DNA packaging: RNA gears in a DNA transport machine. *Cell* **94**:147-150.
75. **Hermoso, J.M., E. Méndez, F. Soriano, and M. Salas.** 1985. Location of the serine residue involved in the linkage between the terminal protein and the DNA of phage ø29. *Nucl. Acids Res.* **13**:7715-7728.
76. **Hoeprich, S., and P. Guo.** 2002. Computer modeling of 3D structure of pRNA monomer, dimer and hexamer of ø29 DNA packaging motor. *J. Biol. Chem.* **277**:20794-20803.
77. **Horcajadas, J.A., W.J.J. Meijer, F. Rojo, and M. Salas.** 2001. Analysis of early promoters of the *Bacillus* bacteriophage GA-1. *J. Bacteriol.* **183**:6965-6970.
78. **Horcajadas, J.A., M. Monsalve, F. Rojo, and M. Salas.** 1999. The switch from early to late transcription in phage GA-1: characterization of the regulatory protein p4_G. *J. Mol. Biol.* **290**:917-928.

79. **Ibarra, B., J.L. Carrascosa, O. Llorca, M. Valle, and J.M. Valpuesta.** 2000. Topology of the components of the DNA packaging machinery in the phage ϕ 29 prohead. *J.Mol.Biol.* **298**:807-815.
80. **Illana, B., L. Blanco, and M. Salas.** 1996. Functional characterization of the genes coding for the terminal protein and DNA polymerase from bacteriophage GA-1. Evidence for a sliding-back mechanism during protein-primed GA-1 DNA replication. *J. Mol. Biol.* **264**:453-464.
81. **Illana, B., J.M. Lázaro, C. Gutiérrez, W.J.J. Meijer, L. Blanco, and M. Salas.** 1999. Phage ϕ 29 terminal protein residues Asn80 and Tyr82 are recognition elements of the replication origins. *J. Biol. Chem.* **274**:15073-15079.
82. **Illana, B., A. Zaballos, L. Blanco, and M. Salas.** 1998. The "RGD" sequence in phage ϕ 29 terminal protein is required for interaction with ϕ 29 DNA polymerase. *Virology* **248**:12-19.
83. **Inciarte, M.R. M. Salas, and J.M. Sogo.** 1980. The structure of replicating DNA molecules of *Bacillus subtilis* phage ϕ 29. *J. Virol.* **34**:187-190.
84. **King, A., W.R. Teertstra, and P.C. van der Vliet.** 1997. Dissociation of the protein primer and DNA polymerase after initiation of adenovirus DNA replication. *J.Biol.Chem.* **272**:24617-24623.
85. **King, A.J, and P.C. van der Vliet.** 1994. A precursor terminal protein-trinucleotide intermediate during initiation of adenovirus DNA replication: regeneration of molecular ends in vitro by a jumping back mechanism. *EMBO J.* **13**:5786-5792.
86. **Kornberg, A., and T.A. Baker.** 1992. DNA replication, W.H. Freeman & Co., New York, N.Y.
87. **Martín, A.C., L. Blanco, P. García, M. Salas, and J. Méndez.** 1996. Protein-primed initiation of pneumococcal phage Cp-1 DNA replication occurs at the third 3' nucleotide of the linear template: a stepwise sliding-back mechanism. *J. Mol. Biol.* **260**:369-377.

88. **Martín, G., J.M. Lázaro, E. Méndez, and M. Salas.** 1989. Characterization of the phage ϕ 29 protein p5 as a single-stranded DNA binding protein. Function in ϕ 29 DNA-protein p3 replication. *Nucl. Acids Res.* **17**:3663-3672.
89. **Matsumoto, K., T. Saito, and H. Hirokawa.** 1983. *In vitro* initiation of bacteriophage ϕ 29 and M2 DNA replication: Genes required for formation of a complex between the terminal protein and 5'dAMP. *Mol.Gen.Genet.* **191**:26-30.
90. **Meijer, W.J.J., J.A. Horcajadas, and M. Salas.** 2001a. The ϕ 29-family of phages. *Microb.Mol.Biol.Rev.* **65**:261-287.
91. **Meijer, W.J.J., P.J. Lewis, J. Errington, and M. Salas.** 2000. Dynamic relocation of phage ϕ 29 DNA during replication and the role of the viral protein p16.7. *EMBO J.* **19**:4182-4190.
92. **Meijer, W.J.J., A. Serna-Rico, and M. Salas.** 2001b. Characterization of the bacteriophage ϕ 29-encoded protein p16.7: a membrane protein involved in phage DNA replication. *Mol. Microbiol.* **39**:731-746.
93. **Mellado, R.P., F. Moreno, E. Viñuela, M. Salas, B.E. Reilly, and D.L. Anderson.** 1976. Genetic analysis of bacteriophage ϕ 29 of *Bacillus subtilis*: Integration and mapping of reference mutants of two collections. *J. Virol.* **19**:495-500.
94. **Mellado, R.P., M.A. Peñalva, M.R. Inciarte, and M. Salas.** 1980. The protein covalently linked to the 5' termini of the DNA of *Bacillus subtilis* phage ϕ 29 is involved in the initiation of DNA replication. *Virology* **104**:84-96.
95. **Mencía, M., M. Monsalve, F. Rojo, and M. Salas.** 1996a. Transcription activation by phage ϕ 29 protein p4 is mediated by interaction with the α subunit of *Bacillus subtilis* RNA polymerase. *Proc. Natl. Acad. Sci. USA* **93**:6616-6620.
96. **Mencía, M., M. Monsalve, M. Salas, and F. Rojo.** 1996b. Transcriptional activator of phage ϕ 29 late promoter: mapping of residues involved in interaction with RNA polymerase and in DNA bending. *Mol. Microbiol.* **20**:273-282.
97. **Méndez, J., L. Blanco, and M. Salas.** 1997. Protein-primed DNA replication: A transition between two modes of priming by a unique DNA polymerase. *EMBO J.* **16**:2519-2527.

98. **Méndez, J., L. Blanco, J.A. Esteban, A. Bernad, and M. Salas.** 1992. Initiation of ϕ 29 DNA replication occurs at the second 3' nucleotide of the linear template: a sliding-back mechanism for protein-primed DNA replication. *Proc. Natl. Acad. Sci. USA* **89**:9579-9583.
99. **Méndez, J., L. Blanco, J.M. Lázaro, and M. Salas.** 1994. Primer-terminus stabilization at the ϕ 29 DNA polymerase active site: mutational analysis of conserved motif Tx₂GR. *J. Biol. Chem.* **269**:30030-30038.
100. **Monsalve, M., B. Calles, M. Mencía, F. Rojo, and M. Salas.** 1998. Binding of phage ϕ 29 protein p4 to the early A2c promoter: recruitment of a repressor by the RNA polymerase. *J. Mol. Biol.* **283**:559-569.
101. **Monsalve, M., B. Calles, M. Mencía, M. Salas, and F. Rojo.** 1997. Transcription activation or repression by phage ϕ 29 protein p4 depends on the strength of the RNA polymerase-promoter interactions. *Molecular Cell* **1**:1-9.
102. **Monsalve, M., M. Mencía, F. Rojo, and M. Salas.** 1995. Transcription regulation in *Bacillus subtilis* phage ϕ 29: expression of the viral promoters throughout the infection cycle. *Virology* **207**:23-31.
103. **Monsalve, M., M. Mencía, F. Rojo, and M. Salas.** 1996a. Activation and repression of transcription at two different phage ϕ 29 promoters are mediated by interaction of the same residues of regulatory protein p4 with the RNA polymerase. *EMBO J.* **15**:101-109.
104. **Monsalve, M., M. Mencía, M. Salas, and F., Rojo.** 1996b. Protein p4 represses phage ϕ 29 A2c promoter by interacting with the α subunit of *Bacillus subtilis* RNA polymerase. *Proc. Natl. Acad. Sci. USA* **93**:8913-8919.
105. **Moreno, F., A. Camacho, E. Viñuela, and M. Salas.** 1974. Suppressor-sensitive mutants of phage ϕ 29. *Virology* **62**:1-16.
106. **Müller, D.J., A. Engel, J.L. Carrascosa, and M. Vélez.** 1997. The bacteriophage ϕ 29 head-tail connector imaged at high resolution with the atomic force microscope in buffer solution. *EMBO J.* **16**:2547-2553.

107. **Nuez, B., and M. Salas.** 1993. Bacteriophage ϕ 29 DNA region controlling late transcription: structural and functional homology with bacteriophage ϕ 29. *Nucl. Acids Res.* **21**:2861-2865.
108. **Nuez, B., F. Rojo, and M., Salas.** 1992. Phage ϕ 29 regulatory protein p4 stabilizes the binding of the RNA polymerase to the late promoter in a process involving direct protein-protein contacts. *Proc. Natl. Acad. Sci. USA* **89**:11401-11405.
109. **Paces, V., C. Vlcek, P. Urbanek, and Z. Hostomsky.** 1986. Nucleotide sequence of the right early region of *Bacillus subtilis* phage PZA completes the 19366-bp sequence of PZA genome. Comparison with the homologous sequence of phage ϕ 29. *Gene* **44**:115-120.
110. **Pastrana, R., J.M. Lázaro, L. Blanco, J.A. García, E. Méndez, and M. Salas.** 1985. Overproduction and purification of protein p6 of *Bacillus subtilis* phage ϕ 29: role in the initiation of DNA replication. *Nucl. Acids Res.* **13**:3083-3100.
111. **Pecenkova, T., and V. Paces.** 1999. Molecular phylogeny of ϕ 29-like phages and their evolutionary relatedness to other protein-primed replicating phages and other phages hosted by Gram-positive bacteria. *J.Mol.Biol.* **48**:197-208.
112. **Pecenkova, T., V. Benes, J. Paces, C. Vlcek, and V. Paces.** 1997. Bacteriophage B103: complete DNA sequence of its genome and relationship to other *Bacillus* phages. *Gene* **199**:157-163.
113. **Peñalva, M.A., and M. Salas.** 1982. Initiation of phage ϕ 29 DNA replication *in vitro*: Formation of a covalent complex between the terminal protein, p3, and 5'-dAMP. *Proc. Natl. Acad. Sci. USA* **79**:5522-5526.
114. **Prieto, I., E. Méndez, and M., Salas.** 1989. Characterization, overproduction and purification of the product of gene 1 of *Bacillus subtilis* phage ϕ 29. *Gene* **77**:195-204.
115. **Prieto, I., M. Serrano, J.M. Lázaro, M. Salas, and J.M. Hermoso.** 1988. Interaction of the bacteriophage ϕ 29 protein p6 with double-stranded DNA. *Proc. Natl. Acad. Sci. USA* **85**:314-318.

116. **Rojo, F., and M. Salas.** 1991. A DNA curvature can substitute phage $\phi 29$ regulatory protein p4 when acting as a transcriptional repressor. *EMBO J.* **10**:3429-3438.
117. **Rojo, F., M. Mencía, M. Monsalve, and M., Salas.** 1998. Transcription activation and repression by interaction of a regulator with the α subunit of RNA polymerase: the model of phage $\phi 29$ protein p4. *Progress Nucleic Acids Res.Mol.Biol.* **60**:29-46.
118. **Saedi, M.S., K.J. Garvey, and J. Ito.** 1987. Cloning and purification of a unique lysozyme produced by Bacillus phage $\phi 29$. *Proc.Natl.Acad.Sci.USA* **84**:955-958.
119. **Salas, M.** 1988. Phages with protein attached to the DNA ends. In *The Bacteriophages*. R. Calendar (ed.) Plenum Publishing Corporation, vol. I, pp. 169-191.
120. **Salas, M.** 1991. Protein-priming of DNA replication. *Annu. Rev. Biochem.* **60**:39-71.
121. **Salas, M.** 1999. Bacillus phage $\phi 29$, in: *Encyclopedia of Virology*, second edition Vol. 1. R.G. Webster and A. Granoff (ed.) pp. 119-130, Saunders Scientific Publications.
122. **Salas, M., J. Méndez, J.A. Esteban, M. Serrano, C. Gutierrez, J.M. Hermoso, A. Bravo, M.S. Soengas, J.M. Lázaro, M.A. Blasco, R. Freire, A. Bernad, J.M. Sogo, and L. Blanco.** 1993. Terminal protein priming of DNA replication: bacteriophage $\phi 29$ as a model system. In *Virus Strategies, Molecular Biology and Pathogenesis*, W. Doerfler and P. Böhm (ed.) Verlag Chemie, pp 3-19.
123. **Salas, M., J.T. Miller, J. Leis, and M.L. DePamphilis.** 1996. Mechanisms for priming DNA synthesis. In *DNA Replication in Eukaryotic Cells*. M. DePamphilis (ed.) Cold Spring Harbor Press, pp. 131-176.
124. **Saturno, J., L. Blanco, M. Salas, and J.A. Esteban.** 1995. A novel kinetic analysis to calculate nucleotide affinity of proofreading DNA polymerases. Application to $\phi 29$ DNA polymerase fidelity mutants. *J. Biol. Chem.* **270**:1-9.
125. **Saturno, J., J.M. Lázaro, L. Blanco, and M. Salas.** 1998. Role of the first aspartate of the "YxDTDS" motif of $\phi 29$ DNA polymerase as a metal ligand during both TP-primed and DNA-primed DNA synthesis. *J. Mol. Biol.* **283**:633-642.

126. **Saturno, J., J.M. Lázaro, F.J. Esteban, L. Blanco, and M. Salas.** 1997. ϕ 29 DNA polymerase residue Lys 383, invariant at motif B of DNA dependent polymerases, is involved in dNTP binding. *J. Mol. Biol.* **269**:313-325.
127. **Serna-Rico, A., B. Illana, M. Salas, and W.J.J. Meijer.** 2000. The putative coiled coil domain of the ϕ 29 terminal protein is a major determinant involved in recognition of the origin of replication. *J. Biol. Chem.* **275**:40529-40538.
128. **Serna-Rico, A., M. Salas, and W.J.J. Meijer.** 2002. The *Bacillus subtilis* phage ϕ 29 protein p16.7, involved in ϕ 29 DNA replication, is a membrane-localized single-stranded DNA-binding protein. *J. Biol. Chem.* **277**:6733-6742.
129. **Serrano, M., C. Gutiérrez, R. Freire, A. Bravo, M. Salas, and J.M. Hermoso.** 1994. Phage ϕ 29 protein p6: a viral histone-like protein. *Biochimie* **76**:981-991.
130. **Serrano, M., C. Gutiérrez, M. Salas, and J.M. Hermoso.** 1993. Superhelical path of the DNA in the nucleoprotein complex that activates the initiation of phage ϕ 29 DNA replication. *J. Mol. Biol.* **230**:248-259.
131. **Serrano, M., J. Gutiérrez, I. Prieto, J.M. Hermoso, and M. Salas.** 1989. Signals at the bacteriophage ϕ 29 DNA replication origins required for protein p6 binding and activity. *EMBO J.* **8**:1879-1885.
132. **Serrano, M., M. Salas, and J.M. Hermoso.** 1990. A novel nucleoprotein complex at a replication origin. *Science* **248**:1012-1016.
133. **Simpson, A.A., Y. Tao, P.G. Leiman, M.O. Badasso, Y. He, P.J. Jardine, N.H. Olson, M.C. Morais, S. Grimes, D.L. Anderson, T.S. Baker, and M.G. Rossmann.** 2000. Structure of the bacteriophage ϕ 29 packaging motor. *Nature* **408**:745-750.
134. **Smith, D.E., S.J. Tans, S.B. Smith, S. Grimes, D.L. Anderson, and C. Bustamante.** 2001. The bacteriophage ϕ 29 portal motor can package DNA against a large internal force. *Nature* **413**:748-752.
135. **Soengas, M.S., J.A. Esteban, J.M. Lázaro, A. Bernad, M.A. Blasco, M. Salas, and L. Blanco.** 1992. Site-directed mutagenesis at the Exo III motif of ϕ 29 DNA polymerase. Overlapping structural domains for the 3'→5' exonuclease and strand-displacement activities. *EMBO J.* **11**:4227-4237.

136. **Soengas, M.S., J.A. Esteban, M. Salas, and C. Gutiérrez.** 1994. Complex formation between phage ϕ 29 single-stranded DNA binding protein and DNA. *J. Mol. Biol.* **239**:213-226.
137. **Soengas, M.S., C. Gutiérrez, and M. Salas.** 1995. Helix-destabilizing activity of ϕ 29 single-stranded DNA binding protein: Effect on the elongation rate during strand displacement DNA replication. *J. Mol. Biol.* **253**:517-529.
138. **Steiner, M., W. Lubitz, and U. Bläsi.** 1993. The missing link in phage lysis of Gram-positive bacteria: gene 14 of bacillus subtilis phage ϕ 29 encodes the functional homolog of lambda S protein. *J. Bacteriol.* **175**:1038-1042.
139. **Talavera, A., F. Jiménez, M. Salas, and E. Viñuela.** 1971. Temperature sensitive mutants of bacteriophage ϕ 29. *Virology* **46**:586-595.
140. **Talavera, A., M. Salas, and E. Viñuela.** 1972. Temperature-sensitive mutants affected in DNA synthesis in phage ϕ 29 of *Bacillus subtilis*. *Eur. J. Biochem.* **31**:367-371.
141. **Tao, Y., N.H. Olson, W. Xu, D.L. Anderson, M.G. Rossmann, and T.S. Baker.** 1998. Assembly of a tailed bacterial virus and its genome release studied in three dimensions. *Cell* **95**:431-437.
142. **Tedin, K., A. Resch, M. Steiner, and U. Bläsi.** 1995. Dual translation start motif evolutionary conserved in the holin gene of *Bacillus subtilis* phage ϕ 29. *Virology* **206**:479-484.
143. **Trottier, M., C.L. Zhang, and P. Guo.** 1996. Complete inhibition of virion assembly in vivo with mutant pRNA essential for phage ϕ 29 DNA packaging. *J. Virol.* **70**:55-61.
144. **Truniger, V., L. Blanco, and M. Salas.** 1999. Role of the "YxGG/A" motif of ϕ 29 DNA polymerase in protein-primed replication. *J. Mol. Biol.* **286**:57-69.
145. **Truniger, V., J.M., Lázaro, M. Salas, and L. Blanco.** 1996. A DNA binding motif coordinating synthesis and degradation in proofreading DNA polymerases. *EMBO J.* **15**:3430-3441.

146. **Truniger, V., J.M. Lázaro, M. Salas, and L. Blanco.** 1998. ϕ 29 DNA polymerase requires the N-terminal domain to bind terminal protein and DNA primer substrates. *J. Mol. Biol.* **278**:741-755.
147. **Valpuesta, J.M., J.M. Carazo, and J.L. Carrascosa.** 1999. The three-dimensional structure of a DNA translocating machine at 10 Å resolution. *Structure* **7**:289-296.
148. **Valle, M., L. Kremer, C. Martínez-A., F. Roncal, J.M. Valpuesta, J.P. Albar, and J.L. Carrascosa.** 1999. Domain architecture of the bacteriophage ϕ 29 connector protein. *J.Mol.Biol.* **288**:899-909.
149. **Vlcek, C., and V. Paces.** 1986. Nucleotide sequence of the late region of *Bacillus* phage ϕ 29 completes the 19285-bp sequence of ϕ 29. Comparison with the homologous sequence of phage PZA. *Gene* **46**:215-225.
150. **Watabe, K., M. Leusch, and J. Ito.** 1984a. Replication of bacteriophage ϕ 29 DNA *in vitro*: The roles of terminal protein and DNA polymerase. *Proc. Natl. Acad. Sci. USA* **81**:5374-5378.
151. **Watabe, K., M. Leusch, and J. Ito.** 1984b. A 3' to 5' exonuclease activity is associated with phage ϕ 29 DNA polymerase. *Biochem.Biophys.Res.Comm.* **123**:1019-1026.
152. **Whiteley, H.R., W.D. Ramey, G.B. Spiegelman, and R.D. Holder.** 1986. Modulation of *in vivo* and *in vitro* transcription of bacteriophage ϕ 29 early genes. *Virology* **155**:392-401.
153. **Yoshikawa, H., K.J. Garvey, and J. Ito.** 1985. Nucleotide sequence analysis of DNA replication origins of the small bacteriophage: evolutionary relationship. *Gene* **37**:125-130.
154. **Zaballos, A. and M. Salas.** 1989. Functional domains in the bacteriophage ϕ 29 terminal protein for interaction with the ϕ 29 DNA polymerase and with DNA. *Nucl. Acids Res.* **17**:10353-10366.
155. **Zhang, F., S. Lemieux, X. Wu, D. St.-Arnaud, T.C. McMurray, F. Major, and D. Anderson.** 1998. Function of hexameric RNA in packaging of bacteriophage ϕ 29 DNA *in vitro*. *Mol.Cell* **2**:141-147.