

## References

1. **Ashbolt, N. J., W. O. K. Grabow, and M. Snozzi.** 2001. Indicators of microbial water quality. In: Water Quality: Guidelines, Standards and Health. L. Fewtrell and J. Bartram., eds. IWA Publishing, London, UK. pp.289-315.
2. **Chung, H., L. A. Jaykus, G. Lovelace, and M. D. Sobsey.** 1998. Bacteriophages and bacteria as indicators of enteric viruses in oysters and their harvest waters. *Water Sci. Technol.* **38**:37-44.
3. **Coetzee, O. J.** 1962. Bakteriophagen als indikator fakaler wassereinigung. *Gesundheits Ingenieur* **12**:371-372.
4. **Dhillon, T. S., Y. S. Chang, S. M. Sun, and W. S. Chau.** 1970. Distribution of coliphages in Hong Kong sewage. *Appl. Microbiol.* **20**:1122-1125.
5. **Dhillon, T. S., E. K. S. Dhillon, H. C. Chau, W. K. Li, and A, H, C. Tsang.** 1976. Studies on bacteriophage distribution - virulent and temperate bacteriophage content of mammalian feces. *Appl. Environ. Microbiol.* **32**:68-74.

6. **Dore, W. J., K. Henshilwood, and D. N. Lees.** 2000. Evaluation of F-specific RNA bacteriophage as a candidate human enteric virus indicator for bivalve molluscan shellfish. *Appl. Environ. Microbiol.* **66**:1280-1295.
7. **Furuse, K.** 1987. Distribution of coliphages in the environment: general considerations. In: S. M. Goyal, C. P. Gerba, and G. Bitton, eds., *Phage Ecology*. Wiley and Sons. NY. pp. 87-124.
8. **Furuse, K., A. Ando, S. Oswa, and I. Wantanabe.** 1981. Distribution of ribonucleic acid coliphages in raw sewage in South and East Asia. *Appl. Environ. Microbiol.* **41**:995-1002.
9. **Gantzer, C., J. Henry, and L. Schwartzbrod.** 2002. *Bacteroides fragilis* and *Escherichia coli* bacteriophages in human feces. *Int. J. Hyg. Environ. Hlth.* **205**:325-328.
10. **Gantzer, C., A. Maul, J. M. Audic, and L. Schwartzbrod.** 1998. Detection of infectious enteroviruses, enterovirus genomes, somatic coliphages, and *Bacteroides fragilis* phages in treated wastewater. *Appl. Environ. Microbiol.* **64**:4307-43012.
11. **Geldreich, E. E.** 1978. Bacterial populations and indicator concepts in feces, sewage, stormwater and soil wastes. In: *Indicators of Viruses in Water and Food*. G. Berg, ed. Ann Arbor Science. Ann Arbor, MI pp 51-98.

12. **Gerba, C. P.** 1987. Phage as indicators of fecal pollution. In: S. M. Goyal, C. P. Gerba, and G. Bitton. *Phage Ecology*. Wiley and Sons, NY. pp.197-210.
13. **Griffin, D. W., R. Stokes, J. B. Rose, and J. H. Paul.** 2000. Bacterial indicator occurrence and the use of an F(+) specific RNA coliphage assay to identify fecal sources in Homosassa Springs, Florida.
14. **Havelaar, A. H.** 1993. A Bacteriophage Standard for Bathing Beaches. final Report. National Institute of Public Health and Environmental Protection. Bilthoven, The Netherlands.
15. **Havelaar, A. H.** 1993b. Bacteriophages as models of human enteric viruses in the environment. *ASM News*. **59**:614-619.
16. **Havelaar, A. H., K. Furuse, and W. M. Hogeboom.** 1986. Bacteriophages and indicator bacteria in human and animal feces. *J. Appl. Bacteriol.* **60**:255-262.
17. **Havelaar, A. H., and W. M. Hogeboom.** 1984. A method for the enumeration of coliphages in sewage and sewage-polluted waters. *Antonie van Leeuwenhoek* **49**:387-397.
18. **Havelaar, A. H., W. M. Hogeboom, and R. Pot.** 1984. F-specific RNA bacteriophages in sewage: methodology and occurrence. *Water Sci. Technol.* **17**:645-655.

Chapter 45, *The Bacteriophages* 2<sup>nd</sup> edition, Calendar & Abedon, Oxford University Press, [thebacteriophages.org](http://thebacteriophages.org)

19. **Havelaar, A. H., M. Olphen, Y. C. Olphen, and Y. C. van Dorst.** 1993. F-specific RNA-bacteriophages are adequate model organisms for enteric viruses in fresh water. *Appl. Environ. Microbiol.* **59**:2956-2962.
20. **Havelaar, A. H., and W. M. Pot-Hogeboom.** 1988. F-specific RNA-bacteriophages as model viruses in water hygiene: ecological aspects. *Water Sci. Technol.* **20**:399-407.
21. **Havelaar, A. H., W. M. Pot-Hogeboom, K. Furuse, R. Pot, and M. P. Hormann.** 1990. F-specific RNA bacteriophages and sensitive host strains in feces and wastewater of human and animal origin. *J. Appl. Bacteriol.* **69**:30-37.
22. **Hsu, F. C., Y. S. Shieh, J. van Duin, M. J. Beekwilder, and M. D. Sobsey.** 1995. Geneotyping male-specific RNA coliphages by hybridization with oligonucleotide probes. *Appl. Environ. Microbiol.* **61**:3960-3966.
23. IAWPRC Study Group on Health Related Water Microbiology. 1991. Bacteriophages as model viruses in water quality control. *Water Res.* **25**:529-545.
24. ISO.1995. International Organization for Standardization. Water Quality. Detection and Enumeration of Bacteriophages-Part 1: Enumeration of F-specific RNA Bacteriophages. ISO/DIS10705-1. Geneva, Switzerland.

Chapter 45, *The Bacteriophages* 2<sup>nd</sup> edition, Calendar & Abedon, Oxford University Press, [thebacteriophages.org](http://thebacteriophages.org)

25. ISO.1999. International Organization for Standardization. Water Quality. Detection and Enumeration of Bacteriophages-Part 2. Enumeration of Bacteriophages Infecting *Bacteroides fragilis*. ISO/DIS10705-4. Geneva, Switzerland.
26. **Jiang, S., R. Noble and W. Chu.** 2001. Human adenoviruses and coliphages in urban runoff-impacted coastal waters of Southern California. *Appl. Environ. Microbiol.* **67**:179-184.
27. **Jofre, J.** 2002. Bacteriophage indicators. In: Encyclopedia of Environmental Microbiology. G. Bitton, ed. John Wiley and Sons, NY. pp 354-363.
28. **Jofre, J., M. Blasi, A. Bosch, and F. Lucena.** 1989. Occurrence of bacteriophages infecting *Bacteroides fragalis* and other viruses in polluted marine sediments. *Water Sci. Technol.* **21**:15-19.
29. **Jofre, J., A. Bosch, R. Lucena, R. Girones, and C. Tartera.** 1986. Evaluation of *Bacteroides fragilis* bacteriophages as indicators of the virological quality of water. *Water Sci. Technol.* **18**:167-173.
30. **Kai, S., S. Wantanabe, K. Furuse and A. Furuse, and A. Osawa.** 1985. *Bacteroides* bacteriophages isolated from human feces. *Microbiol. Immunol.* **29**:895-899.
31. **Kott, Y.** 1966. Estimation of low numbers of *Escherichia coli* bacteriophage by use of the

most probable number method. Appl. Microbiol. **14**:141-144.

**32. Leclerc, H., S. Edberg, V. Pierzo, and J. M. Delattre.** 2000. Bacteriophages as indicators of enteric viruses and public health risk in groundwaters. J. Appl. Microbiol. **88**:5-21.

**33. Osawa, S., K. Furuse and I. Watanabe.** 1981. Distribution of ribonucleic acid coliphages in animals. Appl. Environ. Microbiol. **41**:164-168.

**34. Puig, A., N. Queralt, J. Jofre and R. Araujo.** 1999. Diversity of *Bacteroides fragilis* strains in their capacity to recover phages from human and animal wastes and from fecally polluted wastewater. Appl. Environ. Microbiol. **65**:1772-1776.

**35. Stetler, R. E.** 1984. Coliphages as indicators of enteroviruses. Appl. Environ. Microbiol. **48**:668-670.

**36. Tartera, C., and J. Jofre.** 1987. Bacteriophages active against *Bacteroides fragilis* in sewage-polluted waters. Appl. Environ. Microbiol. **24**:1632-1637.

**37. Woody, M. A., and D. O. Cliver.** 1997. Replication of coliphage Q beta as affected by host cell number, nutrition, competition from insusceptible cells and non-FRNA coliphages. J. Appl. Microbiol. **82**:431-440.